

A CLASS OF GENERALIZED GAMMA FUNCTIONS

JEAN-PAUL JURZAK

ABSTRACT. In this paper, we study the holomorphic function defined by the infinite product $\Gamma_{a,r}(s) = \prod_{n \geq 0} (1 + \frac{1}{a+nr})^s (1 + \frac{s}{a+nr})^{-1}$ which generalize Euler's definition in the sense that $\Gamma(s) = \frac{\Gamma_{1,1}(s)}{s}$. We obtain analogues of the Gauss multiplication formula, complement formula for functions $\Gamma_{a,r}(s)$.

Laboratoire Gevrey de Mathématique Physique, Université de Bourgogne,
Faculté des Sciences et Techniques
BP 47870, 21078 Dijon Cedex
e-mail : Jean-Paul.Jurzak@u-bourgogne.fr

1. INTRODUCTION

Analogues of the gamma function $\Gamma(s)$ have been proposed and studied by L.Bendersky [1], E.L.Post[3], S.Ramanujan [4] and other authors such as F.H.Jackson which introduces a q -gamma function defined as an infinite product.

Euler's original definition of $\Gamma(s)$ is

$$(1.1) \quad \Gamma(s) = \frac{1}{s} \prod_{n \in \mathbb{N}^*} \frac{(1 + \frac{1}{n})^s}{(1 + \frac{s}{n})}$$

and many representations of $\Gamma(s)$ as integrals involving classical functions can be found in classical manuals (see [2] for example).

Our paper differs from previous approaches and study a modified gamma function discribed by formula $\Gamma_{a,r}(s) = \prod_{n \geq 0} (1 + \frac{1}{a+nr})^s (1 + \frac{s}{a+nr})^{-1}$ closed to Euler's definition. However, no simple integral formula (involving classical functions distinct from Euler gamma function) has been obtained as an alternative description of the function $\Gamma_{a,r}(s)$.

The paper is organized as follows: main properties of $\Gamma_{a,r}(s)$ are summed in proposition 3.2 and proofs of results of this section are concentrated in section 4. Section 2 states few properties of infinite products defined by formula 2.1.

2. PRELIMINARIES

For a real sequence $\mathbb{A} = (a_n)_{n \geq 0}$ such that $1 + \frac{1}{a_n} > 0$ for all n , we formally put

$$(2.1) \quad \Gamma_{\mathbb{A}}(s) = \prod_{n \geq 0} \frac{(1 + \frac{1}{a_n})^s}{(1 + \frac{s}{a_n})}$$

Clearly $\Gamma_{\mathbb{A}}(0) = \Gamma_{\mathbb{A}}(1) = 1$.

Definition 2.1. Let $\mathbb{A} = (a_i)_{i \in \mathbb{I}}$ be a subset of $\mathbb{R}$. For $\alpha \in \mathbb{C}^*$ and $n \in \mathbb{Z}$, we define

$$\frac{\mathbb{A}}{\alpha} = (\frac{a_i}{\alpha})_{i \in \mathbb{I}} ; \mathbb{A} + \alpha = (a_i + \alpha)_{i \in \mathbb{I}} ; \mathbb{A}^n = (a_i^n)_{i \in \mathbb{I}}$$

Proposition 2.2. We assume that $\sum_{n=1}^{\infty} \frac{1}{a_n^2} < +\infty$. One has the following properties:

1. The function $\Gamma_{\mathbb{A}}(s)$ is defined and meromorphic on $\mathbb{C}$ with poles only at $s = -a_n$ for $n \in \mathbb{N}$ and never vanishes on $\mathbb{C} - (-\mathbb{A})$.

2. For $\alpha \in \mathbb{R}^*$ and $s \in \mathbb{C} - (-\frac{\mathbb{A}}{\alpha})$, one has:

$$(2.2) \quad \Gamma_{\mathbb{A}}(\alpha s) = \Gamma_{\mathbb{A}}(\alpha)^s \Gamma_{\frac{\mathbb{A}}{\alpha}}(s)$$

3. Assuming $a_n \neq \pm 1$ for all n , one has, for $s \in \mathbb{C} - (\pm \mathbb{A})$

$$(2.3) \quad \Gamma_{\mathbb{A}}(s) = \mu_{\mathbb{A}}^{s-s^2} \frac{\Gamma_{-\mathbb{A}}(s)}{\Gamma_{-\mathbb{A}^2}(s^2)}$$

with $\mu_{\mathbb{A}} = \prod_{n \geq 0} (1 - \frac{1}{a_n^2})$.

In this context, the strictly positive real number $e^{-\gamma_{\mathbb{A}}} = \prod_{n \geq 0} (1 + \frac{1}{a_n}) e^{-\frac{1}{a_n}}$ is well defined and one has, for $s \in \mathbb{C} - (-\mathbb{A})$:

$$(2.4) \quad \frac{\Gamma'_{\mathbb{A}}(s)}{\Gamma_{\mathbb{A}}(s)} = -\gamma_{\mathbb{A}} + \sum_{n=0}^{\infty} \frac{s}{a_n(a_n + s)}$$

Choosing $\mathbb{A} = \mathbb{N}^*$, we get $\Gamma(s) = \frac{1}{s} \Gamma_{\mathbb{N}^*}(s)$ with $\gamma_{\mathbb{A}}$ equal to the Euler constant γ .

Proof:

Putting $J_n = \mathbb{N} \cap [0, n]$ for $n \in \mathbb{N}$, one has, for $s \in \mathbb{R}$:

$$(2.5) \quad \prod_{k \in J_n} \frac{(1 + \frac{1}{a_k})^s}{(1 + \frac{s}{a_k})} = \frac{\prod_{k \in J_n} ((1 + \frac{1}{a_k}) e^{-\frac{1}{a_k}})^s}{\prod_{k \in J_n} (1 + \frac{s}{a_k}) e^{-\frac{s}{a_k}}}$$

The sequence of entire functions $s \rightarrow \prod_{k \in J_n} (1 + \frac{s}{a_k}) e^{-\frac{s}{a_k}}$ is uniformly convergent (as $n \rightarrow +\infty$) on compact sets of $\mathbb{C}$ to the entire function $s \rightarrow \prod_{n \in \mathbb{N}^*} (1 + \frac{s}{a_k}) e^{-\frac{s}{a_k}}$ which vanishes only at points s satisfying $(1 + \frac{s}{a_k}) e^{-\frac{s}{a_k}} = 0$, ie for $s \in \mathbb{A}$. Clearly

$$\prod_{k \in J_n} ((1 + \frac{1}{a_k}) e^{-\frac{1}{a_k}})^s \rightarrow e^{-\gamma_{\mathbb{A}} s}$$

as n tends to $+\infty$, uniformly for s in a compact set of $\mathbb{C}$. Thus, formula 2.1 defines a meromorphic function on $\mathbb{C}$ with poles only at $s = -a_k$ for $n \geq 0$, as a quotient of two entire functions.

For formula 2.2, omitting terms of the form $e^{\pm \frac{1}{a_k}}$, it suffice to prove the equality for $s \in \mathbb{R}^{+*}$ and formula holds for a general $s \in \mathbb{C} - (-\frac{\mathbb{A}}{\alpha})$ by analytic continuation. One has, for $\alpha \in \mathbb{R}^{+*}$

$$\begin{aligned} \frac{\Gamma_{\mathbb{A}}(\alpha s)}{\Gamma_{\mathbb{A}}(\alpha)^s} &= \lim_{n \rightarrow \infty} \prod_{k \in J_n} \frac{(1 + \frac{1}{a_k})^{\alpha s}}{(1 + \frac{\alpha s}{a_k})} \left[\prod_{k \in J_n} \frac{(1 + \frac{\alpha}{a_k})}{(1 + \frac{1}{a_k})^{\alpha}} \right]^s \\ &= \lim_{n \rightarrow \infty} \prod_{k \in J_n} \frac{(1 + \frac{\alpha}{a_k})^s}{(1 + \frac{\alpha s}{a_k})} = \Gamma_{\frac{\mathbb{A}}{\alpha}}(s) \end{aligned}$$

For formula 2.3, one has:

$$\begin{aligned} \Gamma_{\mathbb{A}}(s) &= \prod_{n \geq 0} \frac{(1 + \frac{1}{a_n})^s}{(1 + \frac{s}{a_n})} = \prod_{n \geq 0} \frac{(1 - \frac{1}{a_n^2})^s (1 - \frac{s}{a_n})}{(1 - \frac{1}{a_n})^s (1 - \frac{s^2}{a_n^2})} \\ &= \prod_{n \geq 0} \frac{(1 - \frac{1}{a_n^2})^{s^2} (1 - \frac{s}{a_n}) (1 - \frac{1}{a_n^2})^s}{(1 - \frac{1}{a_n})^s (1 - \frac{s^2}{a_n^2}) (1 - \frac{1}{a_n^2})^{s^2}} = \mu_{\mathbb{A}}^{s-s^2} \frac{\Gamma_{-\mathbb{A}}(s)}{\Gamma_{-\mathbb{A}^2}(s^2)} \end{aligned}$$

We note the

Proposition 2.3. *The functional equation satisfied by the Riemann zeta function is:*

$$(2.6) \quad \frac{\Gamma_{2\mathbb{N}^*}(s) \zeta(s)}{2^s s} = \frac{\Gamma_{2\mathbb{N}^*}(1-s) \zeta(1-s)}{2^{1-s} (1-s)}$$

This equality is the starting point of this work.

Proof:

Since $\Gamma(\frac{1}{2}) = \sqrt{\pi}$, the functional equation takes the form

$$\frac{\Gamma(\frac{s}{2}) \zeta(s)}{\Gamma(\frac{1}{2})^s} = \frac{\Gamma(\frac{1-s}{2}) \zeta(1-s)}{\Gamma(\frac{1}{2})^{1-s}}$$

hence, from $\Gamma(s) = \frac{1}{s} \Gamma_{\mathbb{N}^*}(s)$

$$\frac{\Gamma_{\mathbb{N}^*}(\frac{s}{2}) \zeta(s)}{2^s s \Gamma_{\mathbb{N}^*}(\frac{1}{2})^s} = \frac{\Gamma_{\mathbb{N}^*}(\frac{1-s}{2}) \zeta(1-s)}{2^{1-s} (1-s) \Gamma_{\mathbb{N}^*}(\frac{1}{2})^{1-s}}$$

Taking $\mathbb{A} = \mathbb{N}^*$ and $\alpha = \frac{1}{2}$ in formula 2.2 gives

$$\frac{\Gamma_{\mathbb{N}^*}(\frac{s}{2})}{\Gamma_{\mathbb{N}^*}(\frac{1}{2})^s} = \Gamma_{2\mathbb{N}^*}(s)$$

A direct substitution imply formula 2.6.

3. PROPERTIES OF $\Gamma_{a,r}$ FUNCTIONS

Let $\mathbb{A} = (a + nr)_{n \geq 0} = a + r\mathbb{N}$ be an arithmetical sequence with reason $r > 0$ and first element $a > 0$ (or with reason $r < 0$ and first element $a < -1$). We define

$$(3.1) \quad \Gamma_{a,r}(s) = \prod_{n \geq 0} \frac{(1 + \frac{1}{a+nr})^s}{(1 + \frac{s}{a+nr})}$$

This formula is closed to Euler's definition 1.1 of gamma function expressed as an infinite product. The $\Gamma_{a,r}$ function is expressed by 3.6 in a non-natural way in terms of the well-known Γ function in order to obtain shorter proofs of some properties of $\Gamma_{a,r}$ functions. Some properties of $\Gamma_{a,r}$ are more natural from explicit formula 3.1. We note that $\Gamma(s) = \frac{1}{s} \Gamma_{1,1}(s)$ and formula 2.6 becomes

$$(3.2) \quad \frac{\Gamma_{2,2}(s) \zeta(s)}{2^s s} = \frac{\Gamma_{2,2}(1-s) \zeta(1-s)}{2^{1-s} (1-s)}$$

Following section 2, we put

$$(3.3) \quad \sin_{a,r}(s) = \prod_{n \geq 0} (1 - \frac{s^2}{(a+nr)^2}) = \frac{1}{\Gamma_{a,r}(s) \Gamma_{a,r}(-s)}$$

$$e^{-\gamma_{a,r}} = \prod_{n \geq 0} (1 + \frac{1}{a+nr}) e^{-\frac{1}{a+nr}}$$

Lemma 3.1. *Let $\Psi(s) = \frac{\Gamma'(s)}{\Gamma(s)}$. One has, for $s \in \mathbb{C} - \mathbb{A}$:*

$$(3.4) \quad \sin_{a,r}(s) = \frac{\Gamma^2(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{s}{r}) \Gamma(\frac{a}{r} - \frac{s}{r})}$$

$$(3.5) \quad e^{-\gamma_{a,r}} = \frac{\Gamma(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{1}{r})} e^{\frac{1}{r} \Psi(\frac{a}{r})}$$

$$(3.6) \quad \Gamma_{a,r}(s) = \frac{1}{\Gamma(\frac{a}{r})} \Gamma\left(\frac{s+a}{r}\right) \frac{\Gamma(\frac{a}{r})^s}{\Gamma(\frac{a}{r} + \frac{1}{r})^s}$$

Proposition 3.2. *The function $\Gamma_{a,r}(s)$ has the following properties*

1. For $s \in \mathbb{C} - \mathbb{A}$

$$(3.7) \quad \Gamma_{a,r}(s+r) = \alpha^r \frac{s+a}{r} \Gamma_{a,r}(s)$$

$$\text{with } \alpha = \frac{\Gamma(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{1}{r})}.$$

2. Complement's formula

$$(3.8) \quad \Gamma_{a,r}(s) \Gamma_{a,r}(r-s) = \left(1 - \frac{s}{a}\right) \frac{\Gamma_{a,r}(r)}{\sin_{a,r}(s)}$$

3. Duplication formula

$$(3.9) \quad (2^{2s})^{\frac{1}{r}} \Gamma_{a,r}(s) \Gamma_{a,r}(s + \frac{r}{2}) = \frac{\Gamma_{a,r}(\frac{r}{2})}{\Gamma_{a,r}(a)} \Gamma_{a,r}(2s+a)$$

with

$$\frac{\Gamma_{a,r}(\frac{r}{2})}{\Gamma_{a,r}(a)} = \frac{\sqrt{\pi}}{2^{\frac{2a}{r}-1}} \frac{\Gamma(\frac{a}{r} + \frac{1}{r})^{a-\frac{r}{2}}}{\Gamma(\frac{a}{r})^{1+a-\frac{r}{2}}}$$

4. Multiplication formula

$$n^{\frac{ns}{r}} \prod_{k=0}^{n-1} \Gamma_{a,r}(s + k\frac{r}{n}) = \frac{\Gamma_{a,r}(\frac{r}{n}) \cdots \Gamma_{a,r}(\frac{kr}{n}) \cdots \Gamma_{a,r}(\frac{(n-1)r}{n})}{\Gamma_{a,r}((n-1)a)} \Gamma_{a,r}(ns + (n-1)a)$$

with

$$(3.10) \quad \frac{\Gamma_{a,r}(\frac{r}{n}) \cdots \Gamma_{a,r}(\frac{kr}{n}) \cdots \Gamma_{a,r}(\frac{(n-1)r}{n})}{\Gamma_{a,r}((n-1)a)} = \frac{(2\pi)^{\frac{n-1}{2}}}{n^{\frac{na}{r}-\frac{1}{2}}} \frac{\Gamma(\frac{a}{r} + \frac{1}{r})^{(n-1)(a-\frac{r}{2})}}{\Gamma(\frac{a}{r})^{(n-1)(a+1-\frac{r}{2})}}$$

Thus, the triple formula may be written as

$$\Gamma_{a,r}(3s+2a) = \text{Cte } 3^{\frac{3s}{r}} \Gamma_{a,r}(s) \Gamma_{a,r}(s + \frac{r}{3}) \Gamma_{a,r}(s + \frac{2r}{3})$$

with

$$\text{Cte} = \frac{3^{\frac{3a}{r}-\frac{1}{2}}}{\sqrt{2\pi}} \frac{\Gamma(\frac{a}{r})^{2+2a-r}}{\Gamma(\frac{a}{r} + \frac{1}{r})^{2a-r}} = \frac{\Gamma_{a,r}(2a)}{\Gamma_{a,r}(\frac{r}{3}) \Gamma_{a,r}(\frac{2r}{3})}$$

We put $\Psi_{a,r}(s) = \frac{\Gamma'_{a,r}(s)}{\Gamma_{a,r}(s)}$. One has, for $s \in \mathbb{C} - (a + r\mathbb{N})$:

$$\Psi_{a,r}(s) = \frac{1}{r} \Psi\left(\frac{s+a}{r}\right) + \ln \Gamma\left(\frac{a}{r}\right) - \ln \Gamma\left(\frac{a}{r} + \frac{1}{r}\right)$$

$$\Psi_{a,r}(s+r) - \Psi_{a,r}(s) = \frac{1}{s+a}$$

Various properties are

Proposition 3.3. *Let $\beta(\cdot, \cdot)$ be the usual beta function. We assume $a > 0$ and $r > 0$. With obvious notations, one has:*

1. For $n \in \mathbb{N}^*$

$$(3.11) \quad \Gamma_{a,r}(s) = \Gamma_{a,nr}(s) \Gamma_{a+r,nr}(s) \cdots \Gamma_{a+kr,nr}(s) \cdots \Gamma_{a+(n-1)r,nr}(s)$$

and

$$\frac{\Gamma_{a,nr}(s)}{\Gamma_{a+kr,nr}(s)} = \frac{\beta(\frac{s+a}{nr}; \frac{k}{n})}{\beta(\frac{a}{nr}; \frac{k}{n})} \frac{\beta(\frac{a}{nr}; \frac{k}{n})^s}{\beta(\frac{a+1}{nr}; \frac{k}{n})^s} = \frac{\beta(\frac{s}{nr}; \frac{a+kr}{nr})}{\beta(\frac{s}{nr}; \frac{a}{nr})} \frac{\beta(\frac{a}{nr}; \frac{1}{nr})^s}{\beta(\frac{a+kr}{nr}; \frac{1}{nr})^s}$$

2. For suitable s and h , one has

$$(3.12) \quad \Gamma_{a-h,r}(s+h) = C q^s \Gamma_{a,r}(s)$$

with

$$C = \Gamma(\frac{a}{r}) \frac{\Gamma(\frac{a-h}{r})^{h-1}}{\Gamma(\frac{a-h}{r} + \frac{1}{r})^h} \quad \text{and} \quad q = \frac{\beta(\frac{a-h}{r}; \frac{1}{r})}{\beta(\frac{a}{r}; \frac{1}{r})} = \frac{\beta(\frac{a+1}{r}; \frac{-h}{r})}{\beta(\frac{a}{r}; \frac{-h}{r})}$$

3. One has

$$(3.13) \quad \Gamma_{a,r}(s) = \frac{1}{\Gamma(\frac{a}{r})} \frac{\Gamma_{a,r}(r)^{\frac{s}{r}}}{(\frac{a}{r})^{\frac{s}{r}}} \Gamma(\frac{s}{r} + \frac{a}{r})$$

When $a = r$, the function $\Gamma_{a,a}(s)$ becomes

$$(3.14) \quad \Gamma_{a,a}(s) = \prod_{n \in \mathbb{N}^*} \frac{(1 + \frac{1}{an})^s}{(1 + \frac{s}{an})}$$

with properties

$$(3.15) \quad \sin_{a,a}(s) = a \frac{\sin(\pi \frac{s}{a})}{\pi s}$$

$$(3.16) \quad e^{-\gamma_{a,a}} =_{def} \prod_{n \in \mathbb{N}^*} (1 + \frac{1}{an}) e^{-\frac{1}{an}} = \frac{e^{-\frac{\gamma}{a}}}{\Gamma(1 + \frac{1}{a})} \quad \gamma \text{ Euler constant}$$

$$(3.17) \quad \Gamma_{a,a}(s) = \frac{\Gamma(1 + \frac{s}{a})}{\Gamma(1 + \frac{1}{a})^s}$$

$$(3.18) \quad \Gamma_{a,a}(s+a) = \frac{1}{\Gamma(1 + \frac{1}{a})^a} \frac{s+a}{a} \Gamma_{a,a}(s)$$

$$(3.19) \quad \Gamma_{a,a}(s) \Gamma_{a,a}(a-s) = \frac{1}{\Gamma(1 + \frac{1}{a})^a} (1 - \frac{s}{a}) \frac{\pi \frac{s}{a}}{\sin(\pi \frac{s}{a})}$$

$$(3.20) \quad \Gamma_{a,a}(\frac{a}{b} s) = \Gamma_{a,a}(\frac{a}{b})^s \Gamma_{b,b}(s)$$

$$(3.21) \quad \Psi_{a,a}(s) = \frac{1}{a} \Psi(1 + \frac{s}{a}) - \ln \Gamma(1 + \frac{1}{a})$$

Proposition 3.4. For $\lambda \neq 0$

$$(3.22) \quad \sin_{a,r}(s) = \sin_{\lambda a, \lambda r}(\lambda s) = \sin_{\frac{a}{r}, 1}\left(\frac{s}{r}\right)$$

$$(3.23) \quad \frac{\sin_{a,r}(s+r)}{\left(\frac{s}{r} - \frac{a}{r} + 1\right)} = - \frac{\sin_{a,r}(s)}{\left(\frac{s}{r} + \frac{a}{r}\right)}$$

$$(3.24) \quad \frac{\sin_{a,r}\left(s + \frac{r}{2}\right)}{\left(\frac{s}{r} - \frac{a}{r} + \frac{1}{2}\right)} = - \frac{\Gamma^2\left(\frac{a}{r}\right)}{\Gamma^2\left(\frac{a}{r} + \frac{1}{2}\right)} \sin_{a+\frac{r}{2}, r}(s)$$

$$(3.25) \quad \sin_{a-\frac{r}{2}, r}(s) = \left(1 - \frac{s^2}{\left(a - \frac{r}{2}\right)^2}\right) \sin_{a+\frac{r}{2}, r}(s)$$

For $n \in \mathbb{N}^*$

$$(3.26) \quad \sin_{a,r}(ns) = \sin_{\frac{a}{n}, r}(s) \sin_{\frac{a}{n} + \frac{r}{n}, r}(s) \cdots \sin_{\frac{a}{n} + \frac{(n-1)r}{n}, r}(s)$$

Taking $n = 2$ in formula 3.26, we find

$$(3.27) \quad \sin_{a,r}(2s) = \sin_{\frac{a}{2}, r}(s) \sin_{\frac{a}{2} + \frac{r}{2}, r}(s)$$

Classical descriptions of $\sin(s)$ and $\cos(s)$ as infinite product are equivalent to

$$\frac{\sin(\pi s)}{\pi s} = \sin_{1,1}(s) \quad \text{et} \quad \cos(\pi s) = \sin_{\frac{1}{2}, 1}(s)$$

with the following properties

$$\begin{aligned} \sin_{\frac{1}{2}, 1}(s) &= \pi\left(s + \frac{1}{2}\right) \sin_{1,1}\left(s + \frac{1}{2}\right) \\ \sin_{\frac{1}{2}, 1}(s+1) &= -\sin_{\frac{1}{2}, 1}(s) \end{aligned}$$

due to $\cos(\pi s) = \sin(\pi s + \frac{\pi}{2})$ and periodicity of $\cos(s)$. And $\sin(2\pi s) = 2 \sin(\pi s) \cos(\pi s)$ expresses as

$$\sin_{1,1}(2s) = \sin_{1,1}(s) \sin_{\frac{1}{2}, 1}(s) = \pi\left(s + \frac{1}{2}\right) \sin_{1,1}(s) \sin_{1,1}\left(s + \frac{1}{2}\right)$$

which agree with formulas 3.27 and 3.24 choosing $a = 1$ and $r = 1$.

4. PROOFS

Proof of lemma 3.1:

One has from [2] formula 8.364

$$\prod_{n \geq 0} \left(1 + \frac{w}{a + nr}\right) e^{-\frac{w}{a + nr}} = \frac{\Gamma\left(\frac{a}{r}\right)}{\Gamma\left(\frac{a}{r} + \frac{w}{r}\right)} e^{\frac{w}{r} \Psi\left(\frac{a}{r}\right)}$$

hence $w = 1$ gives formula 3.5. Choosing $w = \pm s$ gives formula 3.4. For $s \in \mathbb{R}$

$$\Gamma_{a,r}(s) = \prod_{n \geq 0} \frac{\left[\left(1 + \frac{1}{a + nr}\right) e^{-\frac{1}{a + nr}}\right]^s}{\left(1 + \frac{s}{a + nr}\right) e^{-\frac{s}{a + nr}}} = \frac{\frac{\Gamma\left(\frac{a}{r}\right)^s}{\Gamma\left(\frac{a}{r} + \frac{1}{r}\right)^s} e^{\frac{s}{r} \Psi\left(\frac{a}{r}\right)}}{\frac{\Gamma\left(\frac{a}{r}\right)}{\Gamma\left(\frac{a}{r} + \frac{s}{r}\right)} e^{\frac{s}{r} \Psi\left(\frac{a}{r}\right)}}$$

thus, by analytic continuation and proposition 2.2

$$\Gamma_{a,r}(s) = \frac{\frac{\Gamma\left(\frac{a}{r}\right)^s}{\Gamma\left(\frac{a}{r} + \frac{1}{r}\right)^s}}{\frac{\Gamma\left(\frac{a}{r}\right)}{\Gamma\left(\frac{a}{r} + \frac{s}{r}\right)}} = \frac{\Gamma\left(\frac{a}{r} + \frac{s}{r}\right)}{\Gamma\left(\frac{a}{r}\right)} \frac{\Gamma\left(\frac{a}{r}\right)^s}{\Gamma\left(\frac{a}{r} + \frac{1}{r}\right)^s}$$

Proof of proposition 3.2:

Let $\alpha = \frac{\Gamma(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{1}{r})}$. One has:

$$\frac{\Gamma_{a,r}(s)}{\Gamma_{a,r}(s+r)} = \frac{1}{\alpha^r} \frac{\Gamma(\frac{s+a}{r})}{\Gamma(\frac{s+r+a}{r})} = \frac{1}{\alpha^r} \frac{\Gamma(\frac{s+a}{r})}{\frac{s+a}{r} \Gamma(\frac{s+a}{r})} = \frac{1}{\alpha^r} \frac{r}{s+a}$$

For complement formula 3.8, with $J_n = \mathbb{N} \cap [0, n]$ for $n \in \mathbb{N}$, one has, for $s \in \mathbb{R}$:

$$\Gamma_{a,r}(s) \Gamma_{a,r}(r-s) = \lim_{n \rightarrow \infty} \prod_{k \in J_n} \frac{1}{1 + \frac{s}{a+kr}} \frac{(1 + \frac{1}{a+kr})^r}{(1 + \frac{r-s}{a+kr})}$$

At fixed n , the denominator is

$$\begin{aligned} & \left(\left(1 + \frac{s}{a}\right) \frac{(a+r+s)}{(a+r)} \frac{(a+2r+s)}{(a+2r)} \dots \frac{(a+nr+s)}{(a+nr)} \right) \left(\frac{(a+r-s)}{a} \frac{(a+2r-s)}{(a+r)} \dots \frac{(a+(n+1)r-s)}{(a+nr)} \right) \\ &= \left(1 + \frac{s}{a}\right) \left[\left(\frac{(a+r)^2 - s^2}{(a+r)^2} \right) \left(\frac{(a+2r)^2 - s^2}{(a+2r)^2} \right) \dots \left(\frac{(a+nr)^2 - s^2}{(a+nr)^2} \right) \right] \frac{(a+(n+1)r-s)}{a} \\ &= \frac{1}{(1 - \frac{s}{a})} \left[\left(1 - \frac{s^2}{a^2}\right) \left(1 - \frac{s^2}{(a+r)^2}\right) \left(1 - \frac{s^2}{(a+2r)^2}\right) \dots \left(1 - \frac{s^2}{(a+nr)^2}\right) \right] \frac{(a+(n+1)r-s)}{a} \end{aligned}$$

Thus

$$\Gamma_{a,r}(s) \Gamma_{a,r}(r-s) = \lim_{n \rightarrow \infty} \left(1 - \frac{s}{a}\right) \frac{a}{(a+(n+1)r-s)} \prod_{k \in J_n} \frac{(1 + \frac{1}{a+kr})^r}{(1 - \frac{s^2}{(a+kr)^2})}$$

One has

$$\begin{aligned} \prod_{k \in J_n} \left(1 + \frac{1}{a+kr}\right)^r &= \prod_{k \in J_n} \left(\left(1 + \frac{1}{a+kr}\right) e^{-\frac{1}{a+kr}} \right)^r e^{\sum_{k \in J_n} \frac{r}{a+kr}} \\ \sum_{k \in J_n} \frac{r}{a+kr} &= \Psi\left(1 + n + \frac{a}{r}\right) - \Psi\left(\frac{a}{r}\right) \end{aligned}$$

Clearly, as $n \rightarrow +\infty$

$$\begin{aligned} \prod_{k \in J_n} \left(1 - \frac{s^2}{(a+kr)^2}\right) &\rightarrow \sin_{a,r}(s) \\ \prod_{k \in J_n} \left(\left(1 + \frac{1}{a+kr}\right) e^{-\frac{1}{a+kr}} \right)^r &\rightarrow e^{-r \gamma_{a,r}} \\ \Psi\left(1 + n + \frac{a}{r}\right) &= \ln\left(1 + n + \frac{a}{r}\right) - \frac{1}{2(1 + n + \frac{a}{r})} + O\left(\frac{1}{n^2}\right) \end{aligned}$$

hence

$$\Gamma_{a,r}(s) \Gamma_{a,r}(r-s) = \left(1 - \frac{s}{a}\right) \frac{e^{-r \gamma_{a,r}}}{\sin_{a,r}(s)} e^{-\Psi(\frac{a}{r})} \lim_{n \rightarrow \infty} \prod_{k \in J_n} \frac{1}{(1 + \frac{(n+1)r-s}{a}) e^{-\Psi(1+n+\frac{a}{r})}}$$

But

$$\prod_{k \in J_n} \left(1 + \frac{(n+1)r-s}{a}\right) e^{-\Psi(1+n+\frac{a}{r})} = \left(1 + \frac{(n+1)r-s}{a}\right) e^{-\ln(1+n+\frac{a}{r}) + \frac{1}{2n} + O(\frac{1}{n^2})} \rightarrow \frac{r}{a}$$

thus

$$\Gamma_{a,r}(s) \Gamma_{a,r}(r-s) = \frac{r}{a} \left(1 - \frac{s}{a}\right) \frac{e^{-r \gamma_{a,r}}}{\sin_{a,r}(s)} e^{-\Psi(\frac{a}{r})}$$

and $s = 0$ gives

$$\Gamma_{a,r}(r) = \frac{r}{a} e^{-r \gamma_{a,r}} e^{-\Psi(\frac{a}{r})}$$

showing 3.8 for $s \in \mathbb{R}$, hence for $s \in \mathbb{C} - (a + r\mathbb{N})$ by analytic continuation.

We prove multiplication formula. Putting $x = \frac{s+a}{r}$ and using formula 3.6, one has

$$\begin{aligned} \prod_{k=0}^{n-1} \Gamma_{a,r}(s + k\frac{r}{n}) &= \prod_{k=0}^{n-1} \frac{1}{\Gamma(\frac{a}{r})} \Gamma(\frac{s+a}{r} + \frac{k}{n}) \frac{\Gamma(\frac{a}{r})^{s+\frac{kr}{n}}}{\Gamma(\frac{a}{r} + \frac{1}{r})^{s+\frac{kr}{n}}} \\ &= \frac{1}{\Gamma(\frac{a}{r})^n} \frac{\Gamma(x)\Gamma(x+\frac{1}{n})\cdots\Gamma(x+\frac{k}{n})\cdots\Gamma(x+\frac{n-1}{n}) \Gamma(\frac{a}{r})^{ns+\frac{(n-1)r}{2}}}{\Gamma(\frac{a}{r} + \frac{1}{r})^{ns+\frac{(n-1)r}{2}}} \end{aligned}$$

From Gauss formula

$$\Gamma(nx) = (2\pi)^{\frac{1-n}{2}} n^{nx-\frac{1}{2}} \prod_{k=0}^{n-1} \Gamma(x + \frac{k}{n})$$

and

$$\Gamma_{a,r}(ns + (n-1)a) \frac{\Gamma(\frac{a}{r} + \frac{1}{r})^{(a+s)n-a}}{\Gamma(\frac{a}{r})^{(a+s)n-a-1}} = \Gamma(nx)$$

we get that

$$\prod_{k=0}^{n-1} \Gamma_{a,r}(s + k\frac{r}{n}) = \frac{\Gamma(\frac{a}{r})^{an-n+a+1+\frac{(n-1)r}{2}}}{(2\pi)^{\frac{1-n}{2}} n^{nx-\frac{1}{2}} \Gamma(\frac{a}{r} + \frac{1}{r})^{a-na+\frac{(n-1)r}{2}}} \Gamma_{a,r}(ns + (n-1)a)$$

thus

$$(2\pi)^{\frac{1-n}{2}} n^{\frac{ns}{r}} n^{\frac{na}{r}-\frac{1}{2}} \prod_{k=0}^{n-1} \Gamma_{a,r}(s + k\frac{r}{n}) = \frac{\Gamma(\frac{a}{r} + \frac{1}{r})^{a-na+\frac{(n-1)r}{2}}}{\Gamma(\frac{a}{r})^{an-n+a+1+\frac{(n-1)r}{2}}} \Gamma_{a,r}(ns + (n-1)a)$$

which agree with 3.10. Taking $n = 2$, we get formula 3.9.

Proof of proposition 3.3:

We prove 1) of proposition 3.3. Formula 3.11 follows from formula 3.1. From formula 3.6, one has

$$\frac{\Gamma_{a,nr}(s)}{\Gamma_{a+kr,nr}(s)} = \frac{\Gamma(\frac{a+kr}{nr})}{\Gamma(\frac{a}{nr})} \frac{\Gamma(\frac{s+a}{nr})}{\Gamma(\frac{s+a+kr}{nr})} \frac{\Gamma(\frac{a}{nr})^s}{\Gamma(\frac{a+kr}{nr})^s} \frac{\Gamma(\frac{a+kr+1}{nr})^s}{\Gamma(\frac{a+1}{nr})^s}$$

Multiplying by $\Gamma(\frac{k}{n})$ or by $\Gamma(\frac{s}{nr})$, one has:

$$\frac{\Gamma(\frac{a+kr}{nr})}{\Gamma(\frac{a}{nr})} \frac{\Gamma(\frac{s+a}{nr})}{\Gamma(\frac{s+a+kr}{nr})} = \frac{\beta(\frac{s+a}{nr}; \frac{k}{n})}{\beta(\frac{a}{nr}; \frac{k}{n})} = \frac{\beta(\frac{s}{nr}; \frac{a+kr}{nr})}{\beta(\frac{s}{nr}; \frac{a}{nr})}$$

In the same way

$$\frac{\Gamma(\frac{a}{nr})^s}{\Gamma(\frac{a+kr}{nr})^s} \frac{\Gamma(\frac{a+kr+1}{nr})^s}{\Gamma(\frac{a+1}{nr})^s} = \frac{\beta(\frac{a}{nr}; \frac{k}{n})^s}{\beta(\frac{a+1}{nr}; \frac{k}{n})^s} = \frac{\beta(\frac{a}{nr}; \frac{1}{nr})^s}{\beta(\frac{a+kr}{nr}; \frac{1}{nr})^s}$$

For 2), one has

$$\frac{\Gamma_{a-h,r}(s+h)}{\Gamma_{a,r}(s)} = C \left[\frac{\Gamma(\frac{a}{r} + \frac{1}{r})}{\Gamma(\frac{a}{r})} \frac{\Gamma(\frac{a-h}{r})}{\Gamma(\frac{a-h}{r} + \frac{1}{r})} \right]^s$$

with

$$C = \Gamma(\frac{a}{r}) \frac{\Gamma(\frac{a-h}{r})^{h-1}}{\Gamma(\frac{a-h}{r} + \frac{1}{r})^h}$$

With a similar method, we get formula 3.12.

For 2), let $\mathbb{A}$ be the sequence $\{a + nr\}_{n \geq 0}$. From 2.2 with $\alpha = r$, we obtain

$$\Gamma_{a,r}(r s) = \Gamma_{a,r}(r)^s \Gamma_{\frac{a}{r},1}(s)$$

From formula 3.6

$$\Gamma_{\frac{a}{r},1}(s) = \frac{1}{\Gamma(\frac{a}{r})} \Gamma(s + \frac{a}{r}) \frac{\Gamma(\frac{a}{r})^s}{\Gamma(\frac{a}{r} + 1)^s} = \frac{1}{(\frac{a}{r})^s \Gamma(\frac{a}{r})} \Gamma(s + \frac{a}{r})$$

thus

$$\Gamma_{a,r}(r s) = \frac{\Gamma_{a,r}(r)^s}{\Gamma(\frac{a}{r})^s (\frac{a}{r})^s} \Gamma(s + \frac{a}{r})$$

which is formula 3.13.

Proof of proposition 3.4:

One clearly has, for $\lambda \in \mathbb{R}^*$, from formula 3.4

$$(4.1) \quad \sin_{a,r}(s) = \frac{\Gamma^2(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{s}{r}) \Gamma(\frac{a}{r} - \frac{s}{r})} = \frac{\Gamma^2(\frac{\lambda a}{\lambda r})}{\Gamma(\frac{\lambda a}{\lambda r} + \frac{\lambda s}{\lambda r}) \Gamma(\frac{\lambda a}{\lambda r} - \frac{\lambda s}{\lambda r})} = \sin_{\lambda a, \lambda r}(\lambda s)$$

Taking $\lambda = \frac{1}{r}$, we get formula 3.22.

For formula 3.23, one has

$$\sin_{a,r}(s + r) = \frac{\Gamma^2(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{s}{r} + 1) \Gamma(\frac{a}{r} - \frac{s}{r} - 1)} = \frac{(\frac{a}{r} - \frac{s}{r} - 1)}{(\frac{a}{r} + \frac{s}{r})} \frac{\Gamma^2(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{s}{r}) \Gamma(\frac{a}{r} - \frac{s}{r})}$$

hence

$$\sin_{a,r}(s + r) = - \frac{(\frac{s}{r} - \frac{a}{r} + 1)}{(\frac{s}{r} + \frac{a}{r})} \sin_{a,r}(s)$$

From formula 3.4, we obtain

$$\sin_{a+\frac{r}{2},r}(s) = \frac{\Gamma^2(\frac{a}{r} + \frac{1}{2})}{\Gamma(\frac{a}{r} + \frac{s}{r} + \frac{1}{2}) \Gamma(\frac{a}{r} - \frac{s}{r} + \frac{1}{2})}$$

thus

$$\sin_{a,r}(s + \frac{r}{2}) = \frac{\Gamma^2(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{s}{r} + \frac{1}{2}) \Gamma(\frac{a}{r} - \frac{s}{r} - \frac{1}{2})} = \frac{(\frac{a}{r} - \frac{s}{r} - \frac{1}{2}) \Gamma^2(\frac{a}{r})}{\Gamma(\frac{a}{r} + \frac{s}{r} + \frac{1}{2}) \Gamma(\frac{a}{r} - \frac{s}{r} + \frac{1}{2})}$$

proving so formula 3.24.

Formula 3.25 follows directly from definition 3.3.

Using formula 3.11, we obtain

$$(4.2) \quad \sin_{a,r}(s) = \sin_{a,nr}(s) \sin_{a+r,nr}(s) \cdots \sin_{a+kr,nr}(s) \cdots \sin_{a+(n-1)r,nr}(s)$$

Replacing s by ns and applying formula 3.22 with $\lambda = \frac{1}{n}$, we obtain formula 3.26.

REFERENCES

- [1] Bendersky, L. Sur la fonction gamma généralisée, *Acta Math.*, 61 (1933), 263-322.
- [2] I.S. Gradshteyn and I.M. Ryzhik *Tables of Integrals, Series, and Products*, 4th ed. Academic Press, New-York, 1965.
- [3] Post, E.L. The generalized gamma function, *Ann. Math.* (2), 20 (1919), 202-217.
- [4] B.C. Berndt *Ramanujan's Notebooks Part I*, Springer-Verlag, New-York, 1985.